

Ovarian Assessment Report

OAR™ with Egg Supply Score (ESS™)

PATIENT: Patient, Example

PHYSICIAN: Clinician Example

DATE OF REPORT: 08/03/2017

PRIVATE AND CONFIDENTIAL

ReproSource®
THE SOURCE FOR FERTILITY TESTING & INFORMATION™

Ovarian Assessment Report 1.5

Patient: Patient, Example

Physician: Clinician Example

Gender: F Age: 28 years Date of Birth: 01/01/1989

Phone: 413-555-5555

Fax: 413-555-5556

Specimen: 00297602

Reported: 08/03/2017

Address: Example Clinic

Received: 07/24/2017

Time: 15:15

123 Example St

Collected: 07/23/2017

Time: 15:15

Example City, EX 55555




TESTS:	FSH	ESTRADIOL	LH	AMH	INHIBIN B
PATIENT VALUES:	8 mIU/mL	30 pg/mL	2.2 mIU/mL	0.51 ng/mL (4% for age)	30 pg/mL
RANGES	FOLLICULAR DAY 2,3 3.0 TO 14.4	ND TO 84	1.1 TO 11.6	Low 0.0 TO <0.3 Reduced 0.3 TO <0.7 Satisfactory 0.7 TO 3.5 High >3.5	0 TO 64 >64
	MIDCYCLE 5.8 TO 21.0	34 TO 400	17.0 TO 77.0	Note: Ranges calibrated to egg supply defined by ovulatory response during egg retrieval ¹ .	
	MENOPAUSAL 21.7 TO 153.0	ND TO 30			

COMMENTS:

Results suggest a reduced chance of a good egg supply. Age suggests a good chance of good egg quality. Given egg supply and quality diminish with time, women experiencing difficulty with conception or considering egg preservation should discuss treatment options with a fertility specialist. Low age specific AMH is associated with premature ovarian insufficiency, earlier menopause, and autoimmunity, while high AMH is associated with PCOS and later menopause. Hormonal contraceptives may lower AMH within the first 8 weeks of use with increasing AMH observed within 8 weeks after discontinuing use. NOTE: Consider repeat testing to confirm concerning serum AMH results. In some women retested within 1 year, AMH can show substantial biological fluctuations. AMH results and ranges are specific to ReproSource. ²⁻⁹

REVIEWED BY: *[Signature]* BENJAMIN LEADER MD PhD

ReproSource is CAP accredited, certified under CLIA '88 to provide high complexity testing, and NY State approved to provide the above testing for clinical use. ReproSource has established, and continues to monitor, the performance characteristics of the above testing. AMH and inhibin B testing has not been cleared or approved by the US FDA and utilizes certain materials labeled "RUO. For Research Use Only." The FDA does not require this testing to go through premarket FDA review. This testing is used for clinical purposes. It should not be regarded as investigational or for research.



UNDERSTANDING YOUR EGG SUPPLY SCORE (ESS)

EGG SUPPLY SCORE CLINICAL RANGES:

Historically, an accurate egg supply assessment was difficult to obtain. The number of eggs obtained through an IVF egg retrieval procedure is considered to be the gold standard for measuring egg supply, but this procedure is not a practical diagnostic test. Although many studies demonstrate the ability of various blood tests to correlate with eggs retrieved, general clinical testing laboratories do not calibrate their test results to this clinical outcome.

The Egg Supply Score (ESS) calibrates age and blood test results from ovary related hormones including AMH and FSH to the number of eggs obtained in an egg retrieval^{10,11}.

The ESS ranges from 1 to 20, with likelihood of good egg supply increasing as the score increases.

ESS	CATEGORY
1 to 5	Reduced
6 to 10	Fair
11 to 15	Good
16 to 20	Excellent

Note: No single test can predict a woman's ability to have a child and the ESS does not assess egg quality.


STUDY CALIBRATING ESS TO EGG SUPPLY:

Objective: To provide the most accurate and easily interpreted assessment of a woman's current egg supply from a blood test.

Methods: In a blinded study^{10,11}, blood samples from women undergoing a total of 454 egg retrievals were tested by ReproSource, and included women with likely excellent egg supply (79 egg donors and 26 female partners of infertile men) and with likely poor egg supply (139 infertile women). ReproSource conducted testing for ovary related hormones, such as AMH and FSH, and for calculation of the ESS. A third party unblinded and analyzed results.

Results: The results (graph to upper right) showed that women with lower ESS values were more likely to have a low egg supply (0 to 4 eggs retrieved, red bars) and women with higher ESS values were more likely to have a good egg supply (12 or more eggs retrieved, dark green bars).

Conclusions: The Egg Supply Score (ESS) is highly correlated with the gold standard measurement of a woman's egg supply: the number of eggs obtained in an egg retrieval procedure. The ESS can be a useful tool to help clinicians and patients better understand a woman's likely egg supply.



AGE SPECIFIC AMH PERCENTILE AND WOMEN'S HEALTH

AMH LEVEL COMPARED TO 65,128 WOMEN* EVALUATED AT FERTILITY CENTERS

HIGH AMH

Condition: Polycystic Ovary Syndrome
IVF/ART: excessive response


AMH
Age Specific Average

Menopause: earlier than average
Conditions: POI/POF, autoimmunity
IVF/ART: poor response
possible decreased egg quality
Exposures: contraceptive hormones, chemotherapy, ovarian surgery

LOW AMH

LEGEND:
Category
Percentile

Borderline High	76% to 85%
Average	26% to 75%
Borderline Low	16% to 25%
Low	6% to 15%
Very Low	0 to 5%



AMH AND WOMEN'S HEALTH

If appropriately calibrated to clinical outcomes, age specific serum AMH testing can provide helpful information for a number of conditions in women's health^{2,3}. Although the population average AMH level declines steadily from 25 years of age until undetectable¹², individual AMH levels vary considerably among women of the same age (graph, above right). The more elevated the age-specific AMH, the higher the likelihood of polycystic ovary syndrome (PCOS)^{6,7} and later menopause^{4,5}. Conversely, lower age-specific AMH values are associated with premature ovarian insufficiency/failure (POI/POF) and earlier menopause. Although not a screening test for autoimmunity, lower serum AMH is observed in conditions such

* **Note:** Women at fertility centers are not randomly selected from the general population but do represent women who are fertile (egg donors), infertile, and randomly selected (partners of infertile men). Thus, percentiles reported may not exactly match the general population.

Graph: Displayed above is the distribution of ReproSource serum AMH values (ng/ml, y-axis) by age (years, x-axis) from 65,128 women evaluated at fertility centers. Color is used to indicate population percentiles, for example, the middle 70% of the population (green) or lowest 5th percentile (red).


as systemic lupus erythematosus and Crohn's disease². Importantly, clinical studies now conclude that hormonal contraceptives can lower AMH values as can ovarian related surgery and chemotherapy^{2,8,9}.

Note: AMH levels are not standardized across laboratories^{2,3}. Thus, AMH results from different laboratories cannot be compared. Secondly, while there is clear clinical benefit from AMH testing, especially as values approach the extremes of high and low, definitive cut points for the general population are still the subject of active research. Therefore, interpretation of AMH results in the general population should be directional rather than definitive, prompting further investigation rather than establishing diagnoses.

WHY FERTILITY EXPERTS USE REPROSOURCE AND THE OVARIAN ASSESSMENT REPORT (OAR)


① Most Up to Date Ovarian Reserve Testing for Clinical Use

ReproSource provides the latest testing related to egg supply and ovarian reserve. Historically FSH has been the blood test most frequently used as a marker of egg supply, but it has a high frequency of falsely reassuring results. AMH, which is secreted from the granulosa cells surrounding each egg, has emerged



as a more accurate blood test^{1,2}. ReproSource has demonstrated AMH is more accurate than other hormones in assessing egg supply¹, declines gradually with age¹², and identifies many women at risk for poor egg supply missed by FSH testing (1 in 11 women tested under age 35 are missed which rises to 1 in 3 women above 39 years of age)¹³.

② Mathematical Formula Calibrates Results to Egg Supply for Easier Interpretation



Clinical reference laboratories report a variety of tests related to egg supply but generally do not calibrate to egg supply¹⁴. Thus, it is often unclear how reported results link to clinical outcomes or how to weight the results of individual tests together. By conducting clinical outcomes research in egg supply testing, ReproSource is able to directly calibrate testing results to egg supply and mathematically optimize the combination of results to provide the Egg Supply Score: a single, easy to use assessment of egg supply.^{10,11}

EXAMPLE PERFORMANCE DETECTING LOW EGG SUPPLY				
TEST	≤4 eggs		≤6 eggs	
	AUROC	vs ESS	AUROC	vs ESS
ESS	0.833	n/a	0.792	n/a
AMH	0.806	P=0.02	0.762	P=0.03
FSH	0.688	P<0.01	0.651	P<0.01

The ESS demonstrated higher accuracy (P<0.05) when compared to single tests such as AMH and FSH for predicting very low (≤4 egg retrieved) or low (≤6 egg retrieved) egg supply when using Area Under the Receiver Operator Characteristic (AUROC) curve comparison^{10,11}.

③ Continuous Improvement of Clinical Information Through Clinical Research

The field of fertility medicine is complex and rapidly evolving with hundreds of studies published each year, many of which rely upon diagnostic testing to categorize patients. Therefore, clinicians need a fertility focused laboratory which calibrates to clinical outcome, and maintains the link between the reported test result and new clinical information¹⁴. ReproSource provides this service.

Recent advancements in egg supply testing, especially related to AMH, have important consequences to women's health. The biggest barrier to the clinical utility of this testing is the frequently changing test methodology, and cut points not calibrated to egg supply^{2,3}. For example, laboratories that provide results for routine tests such as FSH or newer tests such as AMH, generally do not use the same testing methodologies employed in the clinical studies that reported the interpretative ranges of clinical utility. By both conducting clinical outcomes research and providing testing for clinical use, ReproSource provides a reliable source for appropriately calibrated ovarian reserve and other fertility testing used by hundreds of fertility specialists.

REFERENCES (** REPROSOURCE PUBLICATIONS)

1. **Assessment of ovarian reserve with anti-Müllerian hormone: a comparison of the predictive value of anti-Müllerian hormone, follicle-stimulating hormone, inhibin B, and age.** Riggs RM, Duran EH, Baker MW, Kimble TD, Hobeika E, Yin L, et al. *American Journal of Obstetrics and Gynecology*. 2008 Aug;199(2):202 e1-8.
2. **Maximizing clinical utility of antimüllerian hormone (AMH) testing in women's health.** Leader B, Baker VL. *Current Opinions in Obstetrics and Gynecology* 2014 Aug;4, In Press.
3. **A practical approach to recent advances in ovarian reserve testing.** Leader B, Baker V. In: *Biennial Reviews of Infertility* 3rd edition. Editors: Schlegel PN, Fauser BC, Carrell, DT, Racowsky C: Springer, Inc; 2013.
4. **Prediction of age at menopause from assessment of ovarian reserve may be improved by using body mass index and smoking status.** La Marca A, Sighinolfi G, Papaleo E, Cagnacci A, Volpe A, Faddy MJ. *PLOS One*. 2013;8(3):e57005.
5. **The relationship between anti-Müllerian hormone in women receiving fertility assessments and age at menopause in subfertile women: evidence from large population studies.** Dolleman M, Faddy MJ, van Disseldorp J, van der Schouw YT, Messow CM, Leader B, et al. *The Journal of Clinical Endocrinology and Metabolism* 2013 May;98(5):1946-53.
6. **Can anti-Müllerian hormone predict the diagnosis of polycystic ovary syndrome? A systematic review and meta-analysis of extracted data.** Iliodromiti S, Kelsey TW, Anderson RA, Nelson SM. *The Journal of Clinical Endocrinology and Metabolism*. 2013 Aug;98(8):3332-40.
7. **Characterization of women with elevated antimüllerian hormone levels (AMH): correlation of AMH with polycystic ovarian syndrome phenotypes and assisted reproductive technology outcomes.** Tal R, Seifer DB, Khanimov M, Malter HE, Grazi RV, Leader B. *American Journal of Obstetrics and Gynecology*. Epub 2014 Mar 2.
8. **Antimüllerian hormone levels decrease in women using combined contraception independently of administration route.** Kallio S, Puurunen J, Ruokonen A, Vaskivuo T, Piltonen T, Tapanainen JS. *Fertility Sterility*. 2013 Apr;99(5):1305-10.
9. **Comparison of ovarian function markers in users of hormonal contraceptives during the hormone-free interval and subsequent natural early follicular phases.** van den Berg MH, van Dulmen-den Broeder E, Overbeek A, Twisk JW, Schats R, van Leeuwen FE, et al. *Human Reproduction* 2010 Jun;25(6):1520-7.
10. **Index predicts number of oocytes at retrieval in fertile and infertile women.** Leader B, Baca QJ, Stadtmayer L, Riggs R, Rivnay B, L Yin. *Human Reproduction* 2008 Jul; 23(Sup1):i80.
11. **Index outperforms AMH, inhibin B, and FSH in predicting poor egg supply.** Leader B, Quinn E, Sullivan L, Yin L, Riggs R, Stadtmayer L. *Fertility and Sterility* 2008 Sep; 90(1):S263-S264.
12. **Age-specific serum anti-Müllerian hormone values for 17,120 women presenting to fertility centers within the United States.** Seifer DB, Baker VL, Leader B. *Fertility and Sterility* 2011 Feb;95(2):747-50.
13. **High frequency of discordance between antimüllerian hormone and follicle-stimulating hormone levels in serum from estradiol-confirmed days 2 to 4 of the menstrual cycle from 5,354 women in U.S. fertility centers.** Leader B, Hegde A, Baca Q, Stone K, Lannon B, Seifer DB, et al. *Fertility and Sterility* 2012 Oct;98(4):1037-42.
14. **Testing and interpreting measures of ovarian reserve: a committee opinion.** Practice Committee of the American Society for Reproductive Medicine. *Fertility and Sterility* 2012 Dec;98(6):1407-15.

ReproSource®

THE SOURCE FOR FERTILITY TESTING & INFORMATION™

www.reprosource.com tel: 781.937.8893